

seq listing 2AR801
SEQUENCE LISTING

<110> bioMérieux B.V.
Venema, Fokke

<120> Method for lowering both sequence variations and increase of base
line's effects in a diagnostic hybridisation assay, assay for
performing such a method and probe for use in the assay

<130> L/2AR80/MK/57

<150> EP 02 080 125.4
<151> 2002-12-03

<160> 13

<170> PatentIn version 3.2

<210> 1
<211> 26
<212> DNA
<213> Artificial

<220>
<223> Ref

<220>
<221> modified_base
<222> (1)..(26)
<223> I

<220>
<221> misc_feature
<222> (1)..(26)
<223> n = inosine

<400> 1
atcaatgagg angctgcaga ntggga 26

<210> 2
<211> 26
<212> DNA
<213> Artificial

<220>
<223> Me-1

<220>
<221> misc_feature
<222> (1)..(26)
<223> n = inosine; r = 2'-O-Me nucleotide of a

<220>
<221> modified_base
<222> (1)..(26)
<223> 2'-O-methyladenosine

<400> 2
atcrrtgrgg rngctgcrgr ntggga 26

seq listing 2AR801

<210> 3
 <211> 26
 <212> DNA
 <213> Artificial

<220>
 <223> Me-2

<220>
 <221> misc_feature
 <222> (1)..(26)
 <223> n = inosine; r = 2'-O-Me nucleotide of a; k = 2'-O-Me nucleotide
 of g

<220>
 <221> modified_base
 <222> (1)..(26)
 <223> gm

<220>
 <221> modified_base
 <222> (1)..(26)
 <223> I

<220>
 <221> modified_base
 <222> (1)..(26)
 <223> 2'-O-methyladenosine

<400> 3
 atcrrtkakk rnkctkcakr ntkkga

26

<210> 4
 <211> 26
 <212> DNA
 <213> Artificial

<220>
 <223> Me-7

<220>
 <221> misc_feature
 <222> (1)..(26)
 <223> n = inosine; r = 2'-O-Me nucleotide of a; k = 2'-O-Me nucleotide
 of g

<220>
 <221> modified_base
 <222> (1)..(26)
 <223> 2'-O-methyladenosine

<220>
 <221> modified_base
 <222> (1)..(26)
 <223> gm

<220>
 <221> modified_base
 <222> (1)..(26)

seq listing 2AR801

<223> I

<400> 4

atcrrtkakk rnkctkcrkr ntkkka

26

<210> 5

<211> 26

<212> DNA

<213> Artificial

<220>

<223> Ref2

<400> 5

atcaatgagg aagctgcaga atggga

26

<210> 6

<211> 26

<212> DNA

<213> Artificial

<220>

<223> LNA1

<220>

<221> misc_feature

<222> (1)..(26)

<223> b = LNA nucleotide of a; w = LNA nucleotide of t

<400> 6

btcaatgagg aagcwgacaga atggga

26

<210> 7

<211> 26

<212> DNA

<213> Artificial

<220>

<223> LNA2

<220>

<221> misc_feature

<222> (1)..(26)

<223> b = LNA nucleotide of a; w = LNA nucleotide of t

<400> 7

btcaatgagg bagcwgacaga atggga

26

<210> 8

<211> 26

<212> DNA

<213> Artificial

<220>

<223> Ref. molecular beacon

seq listing 2AR801

<220>
 <221> misc_feature
 <222> (1)..(26)
 <223> n = inosine

<220>
 <221> modified_base
 <222> (1)..(26)
 <223> I

<400> 8
 atcaatgagg angctgcaga ntggga

26

<210> 9
 <211> 26
 <212> DNA
 <213> Artificial

<220>
 <223> Subtype A

<220>
 <221> misc_feature
 <222> (1)..(26)
 <223> r = 2'-O-Me nucleotide of a

<220>
 <221> modified_base
 <222> (1)..(26)
 <223> 2'-O-methyladenosine

<400> 9
 atcaatgagg argctgcaga rtggga

26

<210> 10
 <211> 26
 <212> DNA
 <213> Artificial

<220>
 <223> Subtype G1

<220>
 <221> misc_feature
 <222> (1)..(26)
 <223> r = 2'-O-Me nucleotide of a; k = 2'-O-Me nucleotide of g; v =
 2'-O-nucleotide of t

<220>
 <221> modified_base
 <222> (1)..(26)
 <223> 2'-O-methyladenosine

<220>
 <221> modified_base
 <222> (1)..(26)
 <223> gm

<220>

seq listing 2AR801

<221> modified_base
<222> (1)..(26)
<223> 2'-O-methylthymidine

<400> 10
atvaatgarg argctgcaga ktggga

26

<210> 11
<211> 26
<212> DNA
<213> Artificial

<220>
<223> Subtype N

<220>
<221> misc_feature
<222> (1)..(26)
<223> r = 2'-O-Me nucleotide of a; d = 2'-O-Me nucleotide of c

<220>
<221> modified_base
<222> (1)..(26)
<223> 2'-O-methyladenosine

<220>
<221> modified_base
<222> (1)..(26)
<223> cm

<400> 11
atcaatgagg argcrgcaga dtggga

26

<210> 12
<211> 26
<212> DNA
<213> Artificial

<220>
<223> Subtype O1

<220>
<221> misc_feature
<222> (1)..(26)
<223> r = 2'-O-Me nucleotide of a; v = 2'-O-Me nucleotide of t

<220>
<221> modified_base
<222> (1)..(26)
<223> 2'-O-methyladenosine

<220>
<221> modified_base
<222> (1)..(26)
<223> 2'-O-methylthymidine

<400> 12
atcaatgavg argcrgcaga vtggga

26

seq listing 2AR801

<210> 13
<211> 26
<212> DNA
<213> Artificial

<220>
<223> Subtype 03

<220>
<221> misc_feature
<222> (1)..(26)
<223> r = 2'-O-Me nucleotide of a; k = 2'-O-Me nucleotide of g; v =
2'-O-Me nucleotide of t

<220>
<221> modified_base
<222> (1)..(26)
<223> 2'-O-methyladenosine

<220>
<221> modified_base
<222> (1)..(26)
<223> gm

<220>
<221> modified_base
<222> (1)..(26)
<223> 2'-O-methylthymidine

<400> 13
atcaatgagg argckgcaga vtggga

26